DOI: 10.7524/j.issn.0254-6108.2017041901

王森, 袁琪, 韩瑞霞, 等. 环境介质中多溴联苯醚 (PBDEs) 分布特征的研究进展 [J]. 环境化学, 2017, 36(12): 2584-2599.

WANG Sen, YUAN Qi, HAN Ruixia, et al. Distribution of polybrominated diphenyl ethers (PBDEs) in the environment: A review [J]. Environmental Chemistry, 2017, 36(12): 2584-2599.

环境介质中多溴联苯醚 (PBDEs) 分布特征的研究进展*

王森^{1,2**} 袁琪¹ 韩瑞霞² 钱亚利¹ 丁柳屹¹

(1. 西北大学城市与环境学院, 西安, 710127;

2. 中国科学院生态环境研究中心环境化学与生态毒理学国家重点实验室, 北京, 100085)

摘要 由多溴联苯醚 (PBDEs) 所引起的环境问题越来越受到社会各界的关注, 了解环境介质中 PBDEs 的分布特征是对其污染进行有效控制的前提. 本文从非生物介质 (大气、水体、沉积物、污泥及土壤) 和生物体内 (植物、动物、人体及食物链) 两方面入手, 对 PBDEs 在世界各地不同环境介质的分布特征进行综述. 高溴代 PBDEs 因其较低的挥发性和水溶性, 极易吸附于土壤、底泥和空气悬浮颗粒等固体介质, 而低溴代 PBDEs 在各环境介质中都有检出; 生物体中 PBDEs 的分布特征与生物种类、污染源、营养级以及个体组织的特异性有一定关系, 人体血液、母乳、头发等也检出了 PBDEs. 随着北美、欧盟等地区从 2004 年开始逐渐对五溴、八溴和十溴 PBDEs 实施禁用, 其含量在北美和欧盟部分地区有所下降, 但是由于十溴 PBDEs 仍在全球很多地区使用, 又因其具有持久性和长距离迁移性, 要实现环境介质中 PBDEs 含量的控制仍需要一定时间, 因此未来研究仍需对 PBDEs 生产和使用区域开展污染水平及趋势相关研究, 并对 PBDEs 的环境行为深入探讨.

关键词 多溴联苯醚 (PBDEs), 环境介质, 生物体, 分布特征.

Distribution of polybrominated diphenyl ethers (PBDEs) in the environment: A review

WANG Sen^{1,2**} YUAN Qi¹ HAN Ruixia² QIAN Yali¹ DING Liuyi¹

(1. Department of Environmental Sciences, College of Urban and Environmental Sciences, Northwest University, Xi'an, 710127, China; 2. State Key Laboratory of Environmental Chemistry and Ecotoxicology, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing, 100085, China)

Abstract: Environmental problems induced by polybrominated diphenyl ethers (PBDEs) are receiving more and more attention. The knowledge about the distribution of PBDEs is the premise for the effective pollution control. This paper reviewed the distribution of PBDEs in the atmosphere, water, sediments, sludge, soil, plants, animals, human body and food chain all over the world. Owing to the low volatility and water solubility, the higher brominated PBDEs can be easily adsorbed to solid particles like soil, sediment and airborne particles. However, the lower brominated PBDEs are detected in all the environmental media. In living organisms, the distribution characteristics of

2017 年 4 月 19 日收稿 (Received: April 19, 2017).

* 国家自然科学基金 (41303073), 陕西省青年科技新星计划 (2016KJXX-83), 陕西省自然科学基金基础研究计划青年人才项目 (2015JQ4101), 陕西省教育厅科学研究项目 (14JK1731) 和西北大学“优秀青年学术骨干支持计划”资助.

Supported by the National Natural Science Foundation of China Young Scientist Funds (41303073), Shaanxi Province Youth Science and Technology New Star Plan (2016KJXX-83), Young Talent Project of Natural Science Basic Research Plan in Shaanxi Province of China (2015JQ4101), Scientific Research Project of Shaanxi Provincial Department of Education (14JK1731) and Outstanding Youth Academic Backbone Support Program of Northwest University.

** 通讯联系人, Tel: 029-88308427, E-mail: wangsen@nwu.edu.cn

Corresponding author, Tel: 029-88308427, E-mail: wangsen@nwu.edu.cn

PBDEs are depended on biological species, pollution sources, trophic levels and tissue specificity, and PBDEs are also found in human such as blood, breast milk and hair. With the prohibition of penta-, octa-, and deca-BDEs in North American and some regions of European Union, the concentrations of PBDEs in the environment have declined to some degree. However, deca-BDEs is still used in other regions around the world. Due to its persistence and long-distance transport, the control of PBDEs in the environment is still a long-term work. Thus, future research needs to focus on the level, trend, and behavior of PBDEs in the regions where they are produced and used.

Keywords: PBDEs, environmental matrices, living organisms, distribution characteristics.

多溴联苯醚(polybrominated diphenyl ethers, PBDEs)是一种添加型溴代阻燃剂,被广泛应用于塑料、纺织品、电子产品、油漆等领域,因其缺乏化学键的束缚,很容易逸散到大气中,通过大气干湿沉降进入水体、土壤和生物体中,又因其具有难降解性和易于积累的特点,对生态环境构成了巨大危害^[1-2].早在1979年,DeCarlo等报道了美国某PBDEs生产厂家周围的土壤和污泥中BDE-209的存在^[3],两年后在瑞典维斯坎河纺织厂下游鱼体内发现多种PBDEs^[4].Haglund等在波罗的海的斑海豹、灰海豹和鲱鱼体内,健康瑞典男性的脂肪组织,以及商用鱼油等多种样品中检出四、五、六溴代PBDEs^[5].近年来,在偏远地区,比如南极地衣和苔藓样品中也检出12种PBDEs^[6].2003年杨永亮等^[7]在中国青岛近岸沉积物和贻贝中检出21种PBDEs单体,这是国内最早关于PBDEs的报道,随后PBDEs逐渐被报道出现在中国香港海水^[8]、广州大气^[9]等中国其它地区的多种环境介质中.目前PBDEs已在大气、水体、底泥、土壤以及多种生物介质中检出,并广泛分布于世界各地.已有研究证明PBDEs能在动物体内富集,并产生一定程度的健康危害,比如干扰性激素和甲状腺激素^[10-12].由于低剂量的五溴联苯醚(penta-BDEs)和八溴联苯醚(octa-BDEs)就可以引起毒性^[13],从2004年开始欧盟和北美地区逐渐禁止了penta-BDEs、octa-BDEs产品的生产和使用,2006年中国限制了PBDEs的使用,并于2014年开始禁止生产、流通、使用和进出口商用penta、octa-BDEs^[14-15].尽管2008年欧盟和美国的一些州相继采取了禁用十溴联苯醚(deca-BDEs)的措施,但deca-BDEs仍在世界上许多地区使用,同时之前产品中的PBDEs不断释放进入环境,由此所引发的环境问题仍然值得关注^[16].

由于PBDEs的持久性、长距离迁移性及其对生物的毒性,有必要全面认识其在全球范围内各种环境介质中的分布特征.本文总结了近些年来国内外大气、水体(包括沉积物和污泥)、土壤、植物、动物和人体等多种环境介质中PBDEs的分布特征,并对未来的研究方向进行展望,旨在促进PBDEs的控制及管理.

1 环境介质中PBDEs的分布特征(Distribution characteristics of PBDEs in the environmental media)

目前市场上使用最多的商品是deca-BDEs,由于其具有低的挥发性和水溶性,极易吸附于土壤等固体介质上,大部分高溴代PBDEs都沉积在土壤、底泥和空气悬浮颗粒中;而低溴代PBDEs具有相对较高的挥发性和水溶性,在各种环境介质中都有检出.

1.1 大气中PBDEs的分布特征

室温下PBDEs的蒸汽压较低,容易挥发到大气中,并随大气长距离迁移.世界各地大气中都有不同程度PBDEs检出,甚至在北极、青藏高原等背景偏远的地区都有发现^[17-42].PBDEs的蒸汽压随溴代数的增加而降低,因此高溴代PBDEs(如BDE-209)更易与颗粒物结合,而低溴代PBDEs(如BDE-28、47、99)更易在大气中长距离传输^[17].

在背景和偏远地区,长距离传输是其主要来源,PBDEs单体均处于很低的浓度水平,以BDE-47、99为主.在北极,夏季低溴代PBDEs通过挥发进入,冬季PBDEs通过气溶胶从中纬度地区输送到北极^[19],Wang等^[20]研究发现北极 Σ PBDEs(包含BDE-209)的平均浓度($17.3 \text{ pg} \cdot \text{m}^{-3}$)高于北太平洋($12.8 \text{ pg} \cdot \text{m}^{-3}$)及其它背景地区($1.5\text{—}8.5 \text{ pg} \cdot \text{m}^{-3}$),可能是受北美大量使用penta-BDEs及温度效应的影响;且由于大气环流和强烈的空气混合作用,PBDEs在北极地区并无明显的地理分异;而北太平洋地区呈现从中纬度向高纬度地区的递减趋势,可能是由于PBDEs在空气团输送过程中发生了稀释、沉积和

分解等作用.在南极,科学考察站的人类活动是PBDEs的潜在排放源,以BDE-17、28为主, Σ PBDEs(包含BDE-209)($0.7\text{--}3.0\text{ pg}\cdot\text{m}^{-3}$)低于北极^[21].大气中PBDEs浓度存在季节差异,夏季气温升高使PBDEs易于挥发和散逸,冬季降温则使其易于沉降和吸附^[22],如加拿大阿勒特大气中 Σ PBDEs呈现夏季高于冬季的趋势,尤其是低溴代PBDEs的相对百分比^[23].Hung等研究表明北极大气中不同PBDEs单体的倍增时间差异较大(从BDE-209的3.5年到BDE-153的28年),这可能是因为penta-BDEs及octa-BDEs在北美及欧洲一些国家逐步淘汰或禁用,但是由于deca-BDEs产品的持续使用,BDE-209的含量持续上升^[24].

就全球而言,亚洲大气中PBDEs浓度最高,尤其是中国PBDEs(包含BDE-209)的浓度为 $2.0\text{--}787\text{ pg}\cdot\text{m}^{-3}$ ^[25,27-30]普遍高于10年前美国和欧洲($\text{nd}\text{--}85\text{ pg}\cdot\text{m}^{-3}$)^[18,31-34].Deng等^[25]研究发现广州、中国香港的PBDEs浓度分别为 $204\text{--}372\text{ pg}\cdot\text{m}^{-3}$ 和 $33.8\text{--}358\text{ pg}\cdot\text{m}^{-3}$.对于同一区域,大气中PBDEs浓度呈现:城市>乡村^[29]、工业区>非工业区^[26-27]的趋势.Strandberg等研究发现美国芝加哥 Σ PBDEs(包含BDE-209)($33\text{--}77\text{ pg}\cdot\text{m}^{-3}$)远高于乡村($4.4\text{--}21\text{ pg}\cdot\text{m}^{-3}$)^[31].吴辉等^[30]采集了中国某溴代阻燃剂生产地的大气样品,其 Σ PBDEs(包含BDE-209)($16000\text{--}240000\text{ pg}\cdot\text{m}^{-3}$)比广西南宁($31.4\text{--}787\text{ pg}\cdot\text{m}^{-3}$)高2—3个数量级.电子垃圾拆解活动也是大气中PBDEs的重要来源^[35-36],如中国最主要的电子垃圾拆解地之一广东贵屿大气中 Σ PBDEs($5379\text{--}47187\text{ pg}\cdot\text{m}^{-3}$)是广州和中国香港城市区域($33.8\text{--}372\text{ pg}\cdot\text{m}^{-3}$)的58—691倍^[25].

室内空气中PBDEs含量远高于室外,也是室外空气中PBDEs的一个重要来源^[37].室内空气中PBDEs主要来自电器的使用,与室内家具数量、装修程度的关系不明显;由于办公室内电器产品(如电脑、空调等)相对密集和频繁的使用,使其PBDEs浓度通常大于家居环境^[39,43];且无论是办公还是家居环境,同一区域内不同采样点PBDEs分布差异均较大,这可能与电器数量、使用频率以及释放源与采样点距离等因素有关^[37-41,43].Zhu等^[42]研究发现中国东部室内灰尘中 Σ PBDEs($2120\text{ ng}\cdot\text{g}^{-1}$)高于西部($847\text{ ng}\cdot\text{g}^{-1}$),城市高于乡村,表明社会经济发展对其浓度水平影响显著.

1.2 水体、沉积物及污泥中PBDEs的分布特征

大气中的PBDEs可以通过干湿沉降迁移到水体、土壤和生物体等环境介质中,土壤中的PBDEs通过淋溶、径流进入水体,造成水体中PBDEs污染.由于PBDEs具有较高的辛醇-水分配系数($\lg K_{ow}$)和较低的水溶性,水体中PBDEs的含量相对较低^[44],低溴代PBDEs(如BDE-28、47、99)在水中流动性更强,而高溴代PBDEs在污染源附近底泥中的残留量更高^[45].

相关研究表明,仅包含水相的水体中PBDEs以BDE-47和BDE-99为主^[8,46-48],而包含悬浮颗粒相的水域^[49-51]以及河流沉积物和污水处理厂的污泥^[46-74]中,BDE-209含量则相对较高(表1).中国香港、加拿大和瑞士的水体中 Σ PBDEs(不含BDE-209)($\text{nd}\text{--}100.64\text{ pg}\cdot\text{L}^{-1}$)和BDE-209浓度($\text{nd}\text{--}23.2\text{ pg}\cdot\text{L}^{-1}$)均低于美国旧金山湾($15.5\text{--}337\text{ pg}\cdot\text{L}^{-1}$, $12.2\text{--}533\text{ pg}\cdot\text{L}^{-1}$),处于较低污染水平^[8,47,51-52];而珠江口水体中BDE-209浓度极高($76\text{--}5693\text{ pg}\cdot\text{L}^{-1}$),主要来自河流沿岸城市中deca-BDEs的大量使用和排放^[50].由于电子垃圾拆解活动广东清远电子垃圾拆解地水库($23.8\text{--}25.0\text{ ng}\cdot\text{L}^{-1}$)和浙江丰江($22.4\text{ ng}\cdot\text{L}^{-1}$)的表层水体中PBDEs(包含BDE-209)浓度远高于其他地区水体^[53-54].也有研究表明水中PBDEs含量具有明显的季节变化特征,如中国东海表层海水 Σ PBDEs含量在冬季($2.9\text{--}5.5\text{ ng}\cdot\text{L}^{-1}$)高于夏季($0.7\text{--}1.5\text{ ng}\cdot\text{L}^{-1}$)^[55].

BDE-209为沉积物中最主要的单体,比如美国五大湖的沉积物中BDE-209占 Σ PBDEs的83%—98%^[56-58],中国香港红树林的底泥样品中,BDE-209浓度($1.5\text{--}75.9\text{ ng}\cdot\text{g}^{-1}\text{ dw}$)比其他单体高1—2个数量级^[59]. Σ PBDEs(不含BDE-209)在美国五大湖、韩国、加拿大、瑞士、中国长三角、中国南海、中国东海、中国香港、中国台湾等地水平相近($\text{nd}\text{--}14.4\text{ ng}\cdot\text{g}^{-1}\text{ dw}$),而荷兰和珠江三角洲($0.1\text{--}114\text{ ng}\cdot\text{g}^{-1}\text{ dw}$)则比它们高1—2个数量级,珠江三角洲($0.7\text{--}7340\text{ ng}\cdot\text{g}^{-1}\text{ dw}$)和松花江($2.8\text{--}9862\text{ ng}\cdot\text{g}^{-1}\text{ dw}$)的BDE-209浓度远高于其他地区^[56-68].Marvin等研究表明,加拿大尼亚加拉河 Σ PBDEs在1980—1988年间呈持续上升趋势,在1988年之后增速加快并于1995年达到最大值 $35\text{ ng}\cdot\text{g}^{-1}\text{ dw}$ ^[69].天津大沽排污河河口PBDEs和BDE-209浓度均显示上层底泥中含量高于下层底泥,表明近几年我国环境介质中PBDEs含量呈上升趋势^[65].

表 1 PBDEs 在表层水、沉积物和污泥中的分布

Table 1 Distribution of PBDEs in the surface of water, sediments and sludge

国家/地区 Country/Area	年份 Year	PBDEs 单体 PBDEs congeners	Σ PBDEs (均值) ^b Σ PBDEs (mean) ^b	BDE-209 (均值) BDE-209 (mean)	主要单体 Main congeners	文献 Reference
表层水($\text{pg}\cdot\text{L}^{-1}$) ^b						
中国香港沿海	2005	BDE-28,47,99,100,153,156,183	nd—97.8	nd	BDE-28,47,100,183	[8]
美国密歇根湖	2004	BDE-47,66,99,100,153,154	18	na	BDE-47,99	[46]
加拿大温尼伯河	2004	BDE-47,85,99,100,153,154	9.6—100.6	nd	BDE-47,99	[47]
南极表层雪	2011—2012	BDE-17,28,47,66,71,85,99,100,153,154,183	130—340	nd	BDE-47,99	[48]
荷兰	1999	BDE-28,47,99,100,153,209	1.6	0.1—4	BDE-47,99,209	[49]
中国珠江江口	2005	BDE-28,47,99,100,153,154,183,209	26.1—156.9	76—5693	BDE-47,99,209	[50]
瑞士图恩湖	2007	BDE-28,47,99,100,153,183,209	17—78 ^a	12.5—23.2(15.6)	BDE-47,99,209	[51]
中国浙江丰江	2009—2010	BDE-47,99,100,153,154,183,209	22400 ^a	na	BDE-47,153,183,209	[53]
中国清远电子垃圾拆解地	2006	BDE-28,47,66,85,99,100,138,153,154,183,196,197,203,205,206,207,208,209	23800—25000 (24400) ^a	400—410	BDE-28,47,99	[54]
沉积物和污泥($\text{ng}\cdot\text{g}^{-1}\text{dw}$) ^b						
加拿大温尼伯河	2004	BDE-47,85,99,100,153,154,209	1.2—1.6(1.3) ^a	0.5—0.8(0.6)	BDE-47,99,209	[47]
瑞士图恩湖	2005	BDE-28,47,99,100,153,183,209	0.2—5.4 ^a	0.1—5.1(1.0)	BDE-47,99,209	[51]
美国五大湖	2001—2002	BDE-28,47,66,85,99,100,153,154,183,209	0.5—6.3	4—242.0	BDE-47,99,209	[56—58]
中国香港红树林	na	BDE-28,47,99,100,153,154,183,209	0.6—14.4	1.5—75.9	BDE-153,183,209	[59]
中国珠江三角洲	2002—2004	BDE-28,47,66,99,100,138,153,154,183,209	0.1—94.7	0.7—7340	BDE-47,99,209	[63]
中国天津大沽排污河口	2005	BDE-17,28,47,66,71,85,99,100,138,153,154,183,209	0.1—0.5	nd—14.9	BDE-209	[65]
中国台湾西南部河口与沿海	2005	BDE-17,28,47,66,99,100,138,153,154,183,190,205,206,209	nd—1.8	nd—6.3	BDE-209	[66]
中国松花江	2008	BDE-28,47,99,100,153,154,183,209	2.9—9871(397) ^a	2.8—9862	BDE-47,183,209	[67]
中国东海	2013	BDE-15,28,47,99,100,153,154,183,209	0.03—1.3(0.3)	0.2—3.2(0.6)	BDE-47,99,209	[68]
美国加州污水处理厂	na	BDE-47,99,100,153,154,209	1918—2086	1010—1440(1183)	BDE-47,99,209	[70]
德国污水处理厂	2002—2003	BDE-28,47,99,100,153,154,183,209	12.5—288(126)	97.1—2217(429)	BDE-47,99,209	[71]
中国污水处理厂	2005	BDE-17,28,47,66,71,85,99,100,138,153,154,183,209	6.2—57(25.5)	<1—1108.7(68.5)	BDE-47,99,183,209	[72]
科威特污水处理厂	2005—2006	BDE-47,99,100,153,154,183,209	0.9—27.8(9.3)	4.8—1595(181.8)	BDE-47,99,209	[74]
中国珠江三角洲污水处理厂	2006—2007	BDE-28,47,66,85,99,100,153,154,183,196,197,203,205,206,207,208,209	158—23750 ^a	150—22894(6586)	BDE-47,99,209	[76]
捷克污水处理厂	2007	BDE-28,47,49,66,85,99,100,153,154,183,209	205—605	685—1403	BDE-47,99,209	[77]

注:a. Σ PBDEs(含 BDE-209);b. Σ PBDEs(不含 BDE-209);c. 中位数。

Note:a. Σ PBDEs(including BDE-209);b. Σ PBDEs(not including BDE-209);c. median.

污水处理厂污泥中也存在大量 PBDEs,以 BDE-47、99、209 为主^[70]。美国加州污水处理厂的 Σ PBDEs(不含 BDE-209)(1918—2086 $\text{ng}\cdot\text{g}^{-1}\text{dw}$)比其他污水处理厂(0.9—605 $\text{ng}\cdot\text{g}^{-1}\text{dw}$)高 1—2 个数量级,中国珠三角污水处理厂的 BDE-209 浓度(150—22894 $\text{ng}\cdot\text{g}^{-1}\text{dw}$)远高于其他污水处理厂(<1—2217 $\text{ng}\cdot\text{g}^{-1}\text{dw}$)^[70-77]。Wang 等在中国 26 个城市 31 个污水处理厂采样,结果表明其 PBDEs 浓度(不含 BDE-209)(6.2—57 $\text{ng}\cdot\text{g}^{-1}\text{dw}$)低于欧洲和北美国家(12.5—2086 $\text{ng}\cdot\text{g}^{-1}\text{dw}$),BDE-209 平均占 PBDEs 总量的 55%,低于其在沉积物和自然水体中所占比例^[72]。

1.3 土壤中 PBDEs 的分布特征

大气和水体中 PBDEs 可迁移到土壤,其中 PBDEs 的生产、运输及其产品的使用、电子垃圾拆解活动、污水灌溉以及污泥土地利用都是土壤中 PBDEs 的重要来源,同时,土壤中 PBDEs 特别是高溴代 PBDEs 通常被认为来自于大气的干湿沉降^[41],目前世界各地土壤中已检出多种 PBDEs 同系物(表 2)^[26,34,73,78-88].Harrod 等在英国横跨西米兰德的城市乡村带土壤中检出的 Σ PBDEs ($0.07\text{--}3.9\text{ ng}\cdot\text{g}^{-1}\text{ dw}$)^[34]与 Hassanin 等在英国和挪威(欧洲背景)林地和草地土壤中的结果 ($0.07\text{--}12\text{ ng}\cdot\text{g}^{-1}\text{ dw}$)较为一致^[78];而青藏高原的 Σ PBDEs ($0.004\text{--}0.3\text{ ng}\cdot\text{g}^{-1}\text{ dw}$)则远低于其他背景地区土壤^[80,83-85].Cetin 等研究表明土耳其市区 ($8.7\text{--}18.6\text{ ng}\cdot\text{g}^{-1}\text{ dw}$)和工业区 ($0.5\text{--}2840\text{ ng}\cdot\text{g}^{-1}\text{ dw}$)土壤中 Σ PBDEs 高于郊区 ($0.8\text{--}6.8\text{ ng}\cdot\text{g}^{-1}\text{ dw}$)^[26].广东清远电子垃圾拆解地 PBDEs 浓度(包含 BDE-209) ($296\text{ ng}\cdot\text{g}^{-1}\text{ dw}$)比农田 ($15.6\text{--}34.1\text{ ng}\cdot\text{g}^{-1}\text{ dw}$)高 1 个数量级以上^[82].污泥施用增加了土壤中 PBDEs 浓度,最显著的是 BDE-209;西班牙和美国弗吉尼亚的研究表明污泥施用土壤 Σ PBDEs ($30\text{--}689\text{ ng}\cdot\text{g}^{-1}\text{ dw}$, $0.5\text{--}140.5\text{ ng}\cdot\text{g}^{-1}\text{ dw}$)远高于未施污泥的土壤 ($20.7\text{ ng}\cdot\text{g}^{-1}\text{ dw}$, $\text{nd}\text{--}11.0\text{ ng}\cdot\text{g}^{-1}\text{ dw}$)^[73,75].同时,土壤有机质、黏土矿物含量以及海拔高度和土壤深度对 PBDEs 在土壤中的组成和分布有一定影响,如随着土壤深度的增加逐渐降低,低溴代较高溴代 PBDEs 有更强的迁移能力^[85-86];对青藏高原的研究发现,随着海拔的升高 ($1439\text{--}3940\text{ m}$),PBDEs 含量逐渐降低,但上升到 4479 m 时其含量明显增加 ($4500\text{--}4900\text{ m}$)^[83-84].

表 2 PBDEs 在土壤中的分布 ($\text{ng}\cdot\text{g}^{-1}\text{ dw}$)
Table 2 Distribution of PBDEs in soil ($\text{ng}\cdot\text{g}^{-1}\text{ dw}$)

国家/地区 Country/Area	采样地 Sampling sites	年份 Year	PBDEs 单体 PBDEs Congeners	Σ PBDEs (均值) ^a Σ PBDEs (mean) ^a	BDE-209(均值) BDE-209(mean)	主要单体 Main congeners	文献 Reference
英国和挪威 (欧洲背景)	林地和草地	1998	BDE-17, 28, 32, 35, 47, 49, 71, 75, 77, 85, 99, 100, 119, 138, 153, 154, 166, 181, 183, 190	0.07—12(1.4) b	na	BDE-47, 99	[78]
英国	城市乡村带	2003—2004	BDE-28, 47, 99, 100, 153, 154	0.07—3.9 b	na	BDE-47, 99	[34]
西班牙	污泥施用	2005	BDE-47, 99, 100, 153, 154, 183, 196, 206, 207, 208, 209	30—689	24.6—655	BDE-183, 209	[73]
	对照			20.7	14.6	BDE-183, 209	
美国弗吉尼亚	污泥施用	2006	BDE-28, 47, 99, 100, 153, 154, 183, 209	0.5—140.5	nd—78.7	BDE-47, 99, 209	[75]
	对照			nd—11.0	nd—10.0	BDE-47, 99, 209	
中国广东贵屿	酸解地	2004	BDE-3, 7, 15, 28, 33, 47, 49, 66, 71, 77, 85, 99, 100, 119, 126, 138, 153, 154, 183, 190, 197, 203, 207, 209	2720—4250 (3570)	1270	BDE-209	[79]
	打印机垃圾场			893—2890 (1440)	510	BDE-99, 209	
	池塘			263—604(399)	328	BDE-209	
	稻田			34.7—70.9(48.2)	37.3	BDE-209	
	水库			2.0—6.2(3.8)	2.8	BDE-209	
中国青藏高原南部	背景	2005	BDE-17, 28, 47, 66, 71, 85, 99, 100, 138, 153, 154, 183, 190, 209	0.004—0.03(0.01)	nd	BDE-28, 47, 99	[80]
中国东部	电子垃圾拆解地	2007	BDE-28, 47, 66, 85, 99, 100, 138, 153, 154, 209	71.6—5710 (1910)	69.9—5530 (1800)	BDE-47, 99, 209	[81]
	对照城市土壤			0.2—0.4(0.3)	nd—0.3(0.2)	BDE-47, 99, 209	
	农田			0.08—0.3(0.2)	0.09	BDE-47, 99, 209	
	化工园区			2.0—269 (40.6)	1.9—268 (40.3)	BDE-47, 99, 209	
	对照乡村土壤			0.3—0.8 (0.6)	nd—0.8 (0.5)	BDE-47, 99, 209	

注: a. Σ PBDEs(含 BDE-209); b. Σ PBDEs(不含 BDE-209).

Note: a. Σ PBDEs(including BDE-209); b. Σ PBDEs(not including BDE-209).

2 生物体内 PBDEs 的分布特征 (Distribution characteristics of PBDEs in living organism)

环境介质中的 PBDEs 可通过扩散、呼吸、主动和被动吸收等作用进入生物体,并沿着食物链在生物体间迁移富集,最终进入人体,给生态环境和人体健康带来潜在的暴露风险.生物体内 PBDEs 的分布特征与生物种类、所处环境的污染状况以及个体组织的特异性具有一定关系.

2.1 植物中 PBDEs 的分布

目前 PBDEs 已在世界各地的植物体内检出(表 3),包括喜马拉雅山和南极等背景点和偏远地区^[6,89],植物体内的 PBDEs 含量一般在 $\text{pg}\cdot\text{g}^{-1}$ — $\text{ng}\cdot\text{g}^{-1}$ 级^[82,87,89-102].偏远地区的植物体以低溴代 PBDEs 为主,高永飞等^[89]研究发现喜马拉雅山地区沿珠穆朗玛峰北坡海拔 5000 m 以上的茺籽草和棘豆样品中 Σ PBDEs 在 2.1 — $3.6 \text{ ng}\cdot\text{g}^{-1} \text{ lw}$ 之间,属全球背景水平,其主要来源于大气的长距离传输.而在电子垃圾拆解地、PBDEs 生产厂、人口稠密混杂工业区的大都市 PBDEs 含量较高,且大多以高溴代 PBDEs 尤其是 BDE-209 为主^[82,87,91-99].Hu 等^[99]研究发现北京 6 种树树皮中 PBDEs 的均值(包含 BDE-209) ($780 \text{ ng}\cdot\text{g}^{-1} \text{ lw}$) 低于美国阿肯色州 ($5700 \text{ ng}\cdot\text{g}^{-1} \text{ lw}$) 和中国东部电子垃圾拆解地 ($1400 \text{ ng}\cdot\text{g}^{-1} \text{ lw}$),高于欧洲(德国比利时, $72 \text{ ng}\cdot\text{g}^{-1} \text{ lw}$; 意大利里雅斯特, $13 \text{ ng}\cdot\text{g}^{-1} \text{ lw}$) 和韩国 ($140 \text{ ng}\cdot\text{g}^{-1} \text{ lw}$).除乔木外,草本植物的叶片及根系也检出多种 PBDEs,以低溴代 PBDEs 尤其是 BDE-47、99 为主^[100,102].Wang 等^[100]通过分析多种植物根系及叶片中的 Σ PBDEs (3.3 — $94.3 \text{ ng}\cdot\text{g}^{-1} \text{ dw}$),发现植物体低溴代 PBDEs 比例高于土壤.此外,谷物、蔬菜、水果等食物中也检测出了 PBDEs^[90,101].

PBDEs 在不同植物及组织中的分布存在一定差异,同时其含量也随季节变化发生改变.Yogui 等^[6]研究发现 PBDEs 在苔藓体内的含量高于地衣,其差异可能由于植物从大气中吸收 PBDEs 的机制.Hu 等^[99]发现 PBDEs 在不同树种间有着不同的浓度分布特征,以柳树为最高,大的叶表面积使其能更好地与空气中 PBDEs 作用,又因其低脂肪含量,是理想的大气中 PBDEs 被动采样器.St-Amand^[103]发现加拿大渥太华某垃圾填埋场附近云杉针叶中 PBDEs 含量在夏季保持持续上升,而之后到春天呈现下降趋势,春季萌芽期最低,并认为下降是因为冬春交替之际针叶蜡质层的退化和磨损所致.

表 3 PBDEs 在植物体内的分布 ($\text{ng}\cdot\text{g}^{-1} \text{ lw}$)
Table 3 Distribution of PBDEs in plants ($\text{ng}\cdot\text{g}^{-1} \text{ lw}$)

国家/地区 Country/Area	种类 Species	年份 Year	PBDEs 单体 PBDEs congeners	Σ PBDEs(均值) ^b Σ PBDEs(mean) ^b	BDE-209(均值) BDE-209(mean)	主要单体 Main congeners	文献 Reference
南极乔治王岛	地衣、苔藓	2005—2006	BDE-15, 28, 33, 47, 49, 66, 85, 99, 100, 153, 154, 183	0.1—1.2 ^c		BDE-47, 99, 100	[6]
中国喜马拉雅山	茺籽草、镰型棘豆	2005	BDE-17, 28, 47, 66, 71, 99, 100, 153, 183	2.1—3.6(2.9)		BDE-28	[89]
巴基斯坦	小麦、水稻	2013	BDE-28, 35, 47, 100, 99, 154, 153, 183	0.07—46 ^c		BDE-99	[90]
中国莱州湾 BFRs 生产厂	草本植物、超市蔬菜	2006	BDE-28, 47, 99, 100, 153, 154, 183, 206, 207, 208, 209	70—5900	52—3100	BDE-209	[91]
中国 68 个城市	马尾松、杨树等树皮	2004—2007	BDE-17, 28, 47, 49, 66, 71, 85, 99, 100, 138, 153, 154, 183, 190, 209	0.02—48.3(2.8) ^{ac}	37.7—88.4 ^c	BDE-209	[95]
北美	硬木及针叶树树皮	2000—2001	BDE-17, 28, 47, 49, 66, 85, 99, 100, 138, 153, 154, 166, 183, 196, 203, 206, 207, 208, 209	2.3—5700 ^a	nd—5570	BDE-47, 99, 209	[97]
中国北京	松、杨树等树皮	2009—2010	BDE-28, 47, 99, 100, 153, 154, 183, 206, 209	99—3700(780) ^a	95—3600(760)	BDE-209	[99]
西班牙	蔬菜、水果	2006	BDE-47, 99, 100, 153, 154, 183	0.01—0.05 ^d		BDE-47, 99	[101]
纽约和伦敦岛, 北极斯瓦尔巴特群岛	苔藓、丛生虎耳草、仙女木、莎草等	2011—2012	BDE-17, 28, 47, 66, 71, 85, 99, 100, 138, 153, 154, 183, 190	0.04—0.5 ^e		BDE-47, 99, 183	[102]

注: a. Σ PBDEs(含 BDE-209); b. Σ PBDEs(不含 BDE-209); c. dw: 干重; d. ww: 湿重; e. 单位为百分比, 表示占 Σ PBDEs 比例.

Note: a. Σ PBDEs(including BDE-209); b. Σ PBDEs(not including BDE-209); c. dw: Dry weight; d. ww: Wet weight; e. The percentage of the PBDEs content against the Σ PBDEs.

2.2 动物中 PBDEs 的分布特征

对于陆生和水生动物体内 PBDEs 的分布均有大量研究(表 4).陆生动物体内 PBDEs 的分布与其种类、性别及器官的特异性有关,又因其可与污染源直接接触,很多陆生生物都有很高的 PBDEs 含量,且比水生生物更易积累高溴代 PBDEs,以 BDE-209 为最主要单体^[88,104-112].刘娟等^[104]研究发现,公鸡鸡肉和鸡肝组织 Σ PBDEs (42 ng·g⁻¹lw、39 ng·g⁻¹lw) 高于母鸡 (20 ng·g⁻¹lw、9.3 ng·g⁻¹lw),且鸡肉中 PBDEs 含量显著高于鸡肝.Qin 等^[106]研究发现浙江台州电子垃圾拆解区附近的鸡脑中 PBDEs 相比其他组织器官低,血脑屏障保护脑免受 PBDEs 的生物积累,且高溴代 PBDEs 较低溴代更有效.Daso 等^[109]发现南非的地犀鸟和肉垂鹤的蛋壳样品中 PBDEs 分布存在明显的差异,可能是由于这两种鸟类具有不同的食性、栖息地选择、迁移行为以及代谢能力等.Yu 等^[112]研究一个陆生食物网发现 BDE-209 在高营养级生物体内存在脱溴代谢,红隼、麻雀和老鼠体内九溴代 PBDEs (nona-BDEs)/deca-BDEs 的值(1、1.4、1.2)明显大于土壤和草(0.4、0.5).

表 4 PBDEs 在陆生动物体内的分布 (ng·g⁻¹lw)

Table 4 Distribution of PBDEs in terrestrial animals (ng·g⁻¹lw)

国家/地区 Country/Sites	种类 Species	年份 Year	肌肉 Muscle	肝脏 Liver	脑 Brain	脂肪 Fat	Σ PBDEs ^a	主要单体 Main congeners	文献 Reference
中国广东清远电子垃圾回收场	鸡	2005	4.4—4382.9 ^e 3.2—178.6 ^f	3.6—7916.5 ^e 0.7—395.8 ^f			0.7—7916.5	BDE-209	[104]
中国河北白洋淀	鸭子	2005—2006	2.4 0.06 ^d	4.3 0.4 ^d	5.3 0.4 ^d		0.03—0.7 ^{bd}	BDE-47, 99	[105]
中国浙江台州	三黄鸡	2000—2001	1092.4	1077.8	15.2	3138.1	15.2—3138.1	BDE-209	[106]
南非林波波省东部草原	地犀鸟、肉垂鹤 蛋壳	2006					nd—264000 ⁱ 5980—7146000 ^j	BDE-47, 100	[109]
中国喜马拉雅山横断山	大熊猫和 红熊猫	2013	25.3—2045 ^h	38.4—742 ^g 52.6—581 ^h	83.0 ^g	16.4—279 ^h	16.4—2158	BDE-209	[110]
阿拉斯加北部和西部	北极熊	2004—2007		0.2—2.7			0.2—2.7	BDE-47	[111]

注: a. Σ PBDEs (含 BDE-209); b. Σ PBDEs (不含 BDE-209); c. dw: 干重; d. ww: 湿重; e. 公鸡; f. 母鸡; g. 大熊猫; h. 红熊猫; i. 地犀鸟; j. 肉垂鹤.

Note: a. Σ PBDEs (including BDE-209); b. Σ PBDEs (not including BDE-209); c. dw: dry weight; d. ww: wet weight; e. Cock; f. Hen; g. Panda; h. Red panda; i. Ground hornbill; j. Wattled Crane.

虽然整体而言水生生物较陆生生物的 PBDEs 含量偏低,但 PBDEs 已广泛分布于包括偏远地区在内的水生生物体内,并在人类活动较为密集的地区达到较严重污染水平.Yang 等在中国青藏高原高山湖泊的鱼体内检出较低水平的 Σ PBDEs (0.09—4.3 ng·g⁻¹ dw)^[113],而法国罗纳河 4 种鱼体内 Σ PBDEs (4.5—182.5 ng·g⁻¹ dw) 高于欧盟国家的环境质量标准 (42.5×10⁻³ ng·g⁻¹ dw) 4000 倍^[114],泰晤士河等 4 条英国河流中 3 种鱼体内 PBDEs 含量 (2—44 ng·g⁻¹ lw) 也远高于该标准^[115].不同于陆生生物更容易积累高溴代 PBDEs,除了一些污染源 (BDE-209) 附近的地区外,大部分地区的水生生物体内以 BDE-47 为主^[91,113-115].水生生物体内 PBDEs 单体的分布特征也受水生生物种类、组织器官、食物来源和代谢能力等影响^[116-119].Sun 等^[116]研究发现珠江三角洲地区的 3 种鱼体内 PBDEs 同系物分布差别很大,鲮鱼和罗非鱼以 BDE-47 为主,而琵琶鱼以 BDE-209 为主.田胜艳等^[120]研究发现底层鱼比表层鱼更易富集 PBDEs,杂食性鱼类比肉食性鱼类富集的 PBDEs 更高;鱼体内 BDE-47/99 的值高于无脊椎动物,而 BDE-153/154 的值低于无脊椎动物,并认为可能是由于鱼类对 PBDEs 的代谢转化能力更强.

自 1960 年 PBDEs 产品生产以来, PBDEs 逐渐在世界各地的动物体内检出,并随时间基本呈现逐步增加的趋势^[121-125].旧金山湾港海豹体内的 PBDEs 含量在 1989—1998 年间增大了两个数量级 (从 88 ng·g⁻¹ lw 到 8325 ng·g⁻¹ lw)^[121],中国南海雄性江豚体内 PBDEs 含量从 1990 年的 84 ng·g⁻¹ lw 升高到 2001 年的 980 ng·g⁻¹ lw^[122],北极及北大西洋地区鲸、海豹等海洋哺乳动物中 PBDEs 含量在 19 世纪末 20 世纪初达到最大值,反映出全球 PBDEs 工业产品产量和使用量的增加^[123].随着各国相继对 PBDEs 商品的生产和使用采取了一系列限制措施,生物体 PBDEs 含量的增速变缓并开始呈现下降趋

势.北美萨利什海的海豹体内 PBDEs 在 1984—2003 年间呈指数增长,但在 2009 年出现下降,这是因为 2004 年开始禁用 penta-BDEs 和 octa-BDEs^[124]. Houde 等研究发现加拿大北极海豹体内 PBDEs 含量在 1998—2008 年间呈明显增长趋势,随后到 2011 年则持续下降,可能由于 2009 年 penta-BDEs 和 octa-BDEs 的全球监管^[125].

2.3 人体中 PBDEs 的分布特征

关于人体内 PBDEs 分布,主要以母乳、血液和头发为研究对象,除此之外,也有研究在人体其他组织中检出 PBDEs.母乳中 PBDEs 的分布随地区的不同而变化,北美地区远高于其他地区;母乳中 PBDEs 单体以 BDE-47、99、153 为主,且日本、印度尼西亚、韩国、加纳、斯洛伐克、英国、瑞典、希腊、澳大利亚以及美国加州等地的研究都表明 BDE-47 含量最高,然而也有一些亚洲和欧洲国家由于 deca-BDEs 的大量使用,BDE-209 在母乳中有较高含量,甚至成为最主要单体^[126-135]. Park 等研究发现美国加州母乳中 PBDEs 含量(包含 BDE-209)(mean, 134 ng·g⁻¹ lw)超出越南、中国、韩国和日本(mean, 0.4—3.7 ng·g⁻¹ lw)以及欧洲(median, 2—3 ng·g⁻¹ lw) 1 个数量级以上,也高于太平洋西北地区(mean, 93.9、49.3 ng·g⁻¹ lw)、德克萨斯州(6.2—580 ng·g⁻¹ lw)等北美的其他地区^[127]. Meironyté 等发现 1972—1997 年间瑞典母乳中 PBDEs 含量从 0.07 ng·g⁻¹ 上升到 4.0 ng·g⁻¹,所检测到的单体种类也由最初两种(BDE-47, 153)变化为 1996/1997 年的 8 种^[136].

血液中 PBDEs 的分布与人群的饮食习惯、生活习惯、所处区域环境状况及从事的职业有关,个体内也因组织的特异性而发生变化^[137-143]. Fromme 等研究表明 BDE-153 在吸烟者血液中多于不吸烟者^[137]. Uemura 等发现日本普通人群血液中的 PBDEs (2.7—4.8 ng·g⁻¹ lw)以 BDE-209 为主,并认为鱼类的摄入是六溴代以下的 PBDEs 积累的主要因素^[139]. Shen 等研究发现浙江路桥、龙游、天台的儿童血液中 PBDEs 含量和单体组成差异显著,拥有电子垃圾拆解地的路桥污染最严重(32.1 ± 17.5 ng·g⁻¹ lw),以 BDE-99、47、153 为主,龙游和天台污染较轻(12.1 ± 7.6、8.4 ± 4.0 ng·g⁻¹ lw),分别以 BDE-153、156、147 和 BDE-47、116、15 为主^[140]. Zhu 等研究办公室清洁工、大学生、警察 3 类人群的血清 PBDEs 含量,发现办公室清洁工 ΣPBDEs (1.7—1980(140) ng·g⁻¹ lw)远高于大学生(2.1—210(15) ng·g⁻¹ lw)和警察(0.5—150(18) ng·g⁻¹ lw),且以高溴代 PBDEs 为主,这表明清洁工可能通过摄入尘粒而富集更多高溴代 PBDEs^[141]. 屈伟月等研究发现由于胎盘屏障作用婴儿脐血液 ΣPBDEs (1.5—12(3.9) ng·g⁻¹ lw)低于母体血液(1.6—17(4.4) ng·g⁻¹ lw),尤其是高溴代 PBDEs 更难透过胎盘到达胎儿^[142]. Ma 等研究表明美国纽约新生儿血液中 PBDEs 含量在 1997—2002 年间变化不大,随后尤其是 2004—2011 年间出现明显下降^[144].

除母乳和血液外,头发也被用来指示人体内 PBDEs 的分布状况^[96,145-146]. Zheng 等研究表明,PBDEs 在垃圾拆解地周围居民头发内平均含量(43.2 ng·g⁻¹ dw)为普通城市、乡村居民(16.6、10.0 ng·g⁻¹ dw)的 3—4 倍^[146]. 此外,在胎盘组织、精液中也检出 PBDEs^[147-148],中国浙江台州门诊男性患者血液中 ΣPBDEs 约为精液中的两倍,其中血液中 BDE-153、209 含量远高于精液,而 BDE-28、47、99 差异不大,表明低溴代 PBDEs 较高溴代更易迁移到精液^[147].

2.4 食物链生物富集放大作用

大量研究结果表明,陆生和水生食物网均存在 PBDEs 的生物放大,且放大程度依据不同单体种类而变化,BDE-153、154、155 是大多数陆生和水生食物网中生物放大因子(Trophic magnification factors, BMFs)较高的单体^[62,87,149-155]. Nie 等对中国电子垃圾拆解地陆生食物链的研究表明,总 PBDEs 的 BMFs 值为 1.6—3.6,即在所有食物链上存在生物放大作用;而各 PBDEs 单体的 BMFs 值为 <1—11.6,这说明 PBDEs 单体具有不同的生物转化和累积能力^[87]. Shaw 等研究发现大西洋海豹体内的 PBDEs 的 BMFs 值为 17.1—76.5,以 BDE-153 和 BDE-155 最高(148—677, 12—236)^[153]. 生物放大作用还受食物链长度、类型以及生物组织器官的影响. Ma 等研究食物链浮游动物-无脊椎动物-鱼发现 PBDEs 单体的 BMFs 值为 0.61—3.21,而将海鸟纳入该食物链,则该值变为 0.8—4.4^[152]. Zheng 等发现,渤海湾包括浮游动物、无脊椎动物和鱼类在内的 11 种海洋生物体内 6 种 PBDEs 单体的 BMFs 变化范围是 0.2—78.6,最高 BMFs 值为食物链(矛尾复虾虎鱼到脉红螺)中 BDE-154(78.6),各 PBDEs 单体 BMFs 在肌肉和肝脏中也不同^[156]. 在渤海湾食物网的研究发现,随着 PBDEs 单体 lg K_{ow} 的增加,其 TMFs 先增大后减少,当 lg K_{ow} 较

低时 PBDEs 同系物可迅速排出体外,而当 $\lg K_{ow} > 6.87$ 时,生物可利用性降低^[157].吴江平等^[158]等研究发现,大部分生物的 BMFs 值都在 1—30 之间,BDE-154、183、209 的 BMFs 值较其他单体偏低,鸟类的 BMFs 值高于鱼类,哺乳动物 BMFs 值变化较大.

然而也有一些研究发现,随营养级增加生物体内的 PBDEs 含量发生了减少.法国卢瓦尔河和塞纳河食物网的研究表明,营养级越高,PBDEs 含量在一些脂肪含量较高的动物体内(如鳗鱼、尸食性螃蟹)降低,发生“生物稀释”^[159].Kelly 等在研究加拿大北极海洋食物网时发现,PBDEs 的 TMFs 变化范围为 0.7—1.6,只有 BDE-47 超过了 1,BDE-28、66、99、100、118、153、154 均小于 1,可能是由于生物对 PBDEs 的代谢转化和快速排出所致^[93].

3 结论与展望(Conclusion and prospects)

PBDEs 可以随大气、水体和生物长距离迁移,通过大气干湿沉降进入水体、土壤和生物体,并在不同环境介质间发生迁移,目前在世界各地多种环境介质中均有检出.关于环境介质中 PBDEs 的时空分布特征的研究全面而广泛,不同环境介质的研究现状为:大气的相关研究多于水体和土壤,而水体又以沉积物居多;对于生物的研究,陆生植物多于水生植物,水生动物多于陆生动物;人体则以直接暴露或者是污染源附近的居民等特殊暴露人群作为主要研究对象,而普通人群的研究较少,其中血液、母乳及头发是研究最多的介质.全球各区域间研究现状也存在较大差异:北美和欧洲的研究起步较早,而非洲、南美洲、亚洲的部分地区研究较少,不利于从全球尺度把握 PBDEs 的污染现状,我国虽然起步较晚,但近些年进展迅速.PBDEs 的分布特征表现为:偏远地区的大气中主要以低溴代 PBDEs 为主,同时低溴代 PBDEs 也是水体的主要单体,而高溴代 PBDEs 则在大气悬浮颗粒物、污染源附近的底泥、土壤以及污水处理厂的污泥中有较高的百分含量.生物体中 PBDEs 的分布特征与生物种类、污染源、营养级以及个体组织的特异性有一定关系,如陆生生物含量普遍高于水生生物,高营养级生物大都高于低营养级生物;生物种属间和组织间的差异可能是由于其吸收、转化能力不同.全球各区域对商品化 PBDEs 生产和使用的不同带来其全球时空分布的差异:北美以 penta-、octa-BDEs 产品为主,其环境中主要单体多为 penta-、octa-BDEs;而亚洲和欧洲更多使用 deca-BDEs,通常 deca-BDEs 是最主要单体.总体来说,20 世纪之前环境中 PBDEs 含量呈上升趋势,2004 年开始各国陆续对 penta-BDEs、octa-BDEs 产品的生产和使用采取一些措施后,其增长率开始变小,部分区域的含量呈现下降趋势,尤其是在北美和欧洲.然而在亚洲等很多地区 deca-BDEs 仍在使用,导致其含量仍持续增加.通过对不同环境介质中 PBDEs 时空分布特征的分析,能够进一步了解全球各地各圈层的 PBDEs 污染源、分布特征,为对比分析国内外污染现状和水平提供数据基础,为 PBDEs 污染控制法律法规政策的制定提供参考.

由于各地 PBDEs 商品的流通以及 PBDEs 的持久性和长距离迁移性,实现环境中 PBDEs 的控制仍需很长时间,因此未来需重点关注仍在使用 deca-BDEs 产品的区域,以及 PBDEs 的生产加工地和电子垃圾拆解地等典型污染区域;并增加对于现有研究较少的地区及研究对象的研究;比如从特殊人群扩大到普通大众,农作物和加工食品中的分布也需更加详细地了解;PBDEs 在各环境介质间会发生一定的迁移和转化,未来需深入探讨其在大气、土壤、水体和生物体中的迁移转化机制,以进一步分析其归趋.这些研究工作将有利于全球尺度把握 PBDEs 的污染现状和趋势,了解其在不同地区的时空分布特点,从而更加有针对性地对污染进行控制.

参考文献(References)

- [1] 刘汉霞,张庆华,江桂斌.多溴联苯醚及其环境问题[J].化学进展,2005,17(3):554-562.
LIU H X, ZHANG Q H, JIANG G B. Polybrominated diphenyl ethers and its related environmental problems[J]. Prgress in Chemistry, 2005, 17(3):554-562(in Chinese).
- [2] MARTIN M, LAM P K, RICHARDSON B J. An Asian quandary: Where have all of the PBDEs gone? [J]. Marine Pollution Bulletin, 2004, 49(5-6):375-382.
- [3] DECARLO V J. Studies on brominated chemicals in the environment[J]. Annals New York Academy of Sciences, 1979, 320(1):678-681.
- [4] ANDERSSON O, BLOMKVIST G. Polybrominated aromatic pollutants found in fish in Sweden[J]. Chemosphere, 1981, 10(9):1051-1060.
- [5] HAGLUND P S, ZOOK D R, BUSER H R, et al. Identification and quantification of polybrominated diphenyl ethers and methoxy-

- polybrominated diphenyl ethers in Baltic biota[J]. *Environmental Science & Technology*, 1997, 31(11):3281-3287.
- [6] YOGUI G T, SERICANO J L. Polybrominated diphenyl ether flame retardants in lichens and mosses from King George Island, maritime Antarctica[J]. *Chemosphere*, 2008, 73(10):1589-1593.
- [7] 杨永亮,潘静,李悦,等. 青岛近岸沉积物中持久性有机污染物多氯萘和多溴联苯醚[J]. *科学通报*, 2003, 48(21):2244-2251.
YANG Y L, PAN J, LI Y, et al. Persistent organic pollutants PCNs and PBDEs in sediments from costal waters of Qingdao, Shandong Peninsula[J]. *Chinese Science Bulletin*, 2003, 48(21):2244-2251 (in Chinese).
- [8] WURL O, LAM P K S, OBBARD J P. Occurrence and distribution of polybrominated diphenyl ethers (PBDEs) in the dissolved and suspended phases of the sea-surface microlayer and seawater in Hong Kong, China[J]. *Chemosphere*, 2006, 65(9):1660-1666.
- [9] CHEN L G, MAI B X, BI X H. Concentration levels, compositional profiles, and gas-particle partitioning of polybrominated diphenyl ethers in the atmosphere of an urban city in South China[J]. *Environmental Science & Technology*, 2006, 40(4):1190-1196.
- [10] 韦朝海,廖建波,刘浔,等. PBDEs的来源特征、环境分布及污染控制[J]. *环境科学学报*, 2015, 35(10):3025-3041.
WEI C H, LIAO J B, LIU X, et al. Source, characteristics, environmental distribution and pollution control of PBDEs[J]. *Acta Scientiae Circumstantiae*, 2015, 35(10):3025-3041 (in Chinese).
- [11] DINGEMANS M M, VAN DEN BERG M, WESTERINK R H. Neurotoxicity of brominated flame retardants: (in) direct effects of parent and hydroxylated polybrominated diphenyl ethers on the (developing) nervous system[J]. *Environmental Health Perspectives*, 2011, 119(7):900-907.
- [12] KODAVANTI P R, COBUM C G, MOSER V C. Developmental exposure to a commercial PBDE mixture, DE-71: Neurobehavioral, hormonal, and reproductive effects[J]. *Toxicological Sciences*, 2010, 116(1):297-312.
- [13] DAMERUD P O. Toxic effects of brominated flame retardants in man and in wildlife[J]. *Environment International*, 2003, 29(6):841-853.
- [14] 李子扬,陈永亨. 多溴联苯醚的环境行为及其生态毒理效应[J]. *科学技术与工程*, 2011, 11(1):97-105.
LI Z Y, CHEN Y H. The environmental behavior and ecotoxicological effects of polybrominated diphenyl ethers (PBDEs) [J]. *Science Technology and Engineering*, 2011, 11(1):97-105 (in Chinese).
- [15] 赵静,徐挺,白建峰. 多溴联苯醚暴露的神经行为效应及其毒理机制[J]. *生态毒理学报*, 2017, 12(1):51-61.
ZHAO J, XU T, BAI J F. The neurobehavioral toxicity induced by polybrominated diphenyl ethers exposure and the underlying mechanisms [J]. *Asian Journal of Ecotoxicology*, 2017, 12(1):51-61 (in Chinese).
- [16] BETTS K. Does a key PBDE break down in the environment? [J]. *Environmental Science & Technology*, 2008, 42(18):6781.
- [17] TITTELMIER S A, HALLDORSON T, STERN G A, et al. Vapor pressures, aqueous solubilities, and Henry's law constants of some brominated flame retardant[J]. *Environmental Toxicology and Chemistry*, 2002, 21(9):1804-1810.
- [18] LEE R G, THOMAS G O, JONES K C. PBDEs in the atmosphere of three locations in western Europe[J]. *Environmental Science & Technology*, 2004, 38(3):699-706.
- [19] SU Y S, HUNG H, SVERKO E, et al. Multi-year measurements of polybrominated diphenyl ethers (PBDEs) in the Arctic atmosphere[J]. *Atmospheric Environment*, 2007, 41(38):8725-8735.
- [20] WANG X M, DING X, MAI B X, et al. Polybrominated diphenyl ethers in airborne particulates collected during a research expedition from the Bohai Sea to the Arctic[J]. *Environmental Science & Technology*, 2005, 39(20):7803-7809.
- [21] LI Y M, GENG D W, LIU F B, et al. Study of PCBs and PBDEs in King George Island, Antarctica, using PUF passive air sampling[J]. *Atmospheric Environment*, 2012, 51(5):140-145.
- [22] 吴库生,刘俊晓,李燕,等. 多溴联苯醚的环境分布[J]. *职业与健康*, 2008, 24(22):2467-2469.
WU K S, LIU J X, LI Y, et al. Environmental distribution of polybrominated diphenyl ethers[J]. *Occup and Health*, 2008, 24(22):2467-2469 (in Chinese).
- [23] XIAO H, SHEN L, SU Y S, et al. Atmospheric concentrations of halogenated flame retardants at two remote locations: The Canadian High Arctic and the Tibetan Plateau[J]. *Environmental Pollution*, 2012, 161(1):154-161.
- [24] HUNG H, KALLENBORN R, BREIVIK K, et al. Atmospheric monitoring of organic pollutants in the Arctic under the Arctic Monitoring and Assessment Programme (AMAP): 1993—2006[J]. *Science of the Total Environment*, 2010, 408(15):2854-2873.
- [25] DENG W J, ZHENG J S, BI X H, et al. Distribution of PBDEs in air particles from an electronic waste recycling site compared with Guangzhou and Hong Kong, South China[J]. *Environment International*, 2007, 33(8):1063-1069.
- [26] CETIN B, ODABASI M. Particle-phase dry deposition and air-soil gas-exchange of polybrominated diphenyl ethers (PBDEs) in Izmir, Turkey[J]. *Environmental Science & Technology*, 2007, 41(14):4986-4992.
- [27] YU Z Q, LIAO R, LI H, et al. Partical-bound dechlorane plus and polybrominated diphenyl ethers in ambient air around Shanghai, China [J]. *Environmental Pollution*, 2011, 159(10):2982-2988.
- [28] QI H, LI W L, LIU L Y, et al. Brominated flame retardants in the urban atmosphere of Northeast China: Concentrations, temperature dependence and gas-particle partitioning[J]. *Science of the Total Environment*, 2014, 491-492:60-66.
- [29] HE W, QIN N, HE Q S, et al. Atmospheric PBDEs at rural and urban sites in central China from 2010 to 2013: Residual levels, potential sources and human exposure[J]. *Environmental Pollution*, 2014, 192:232-243.

- [30] 吴辉, 金军, 王英, 等. 典型地区大气中多溴联苯醚和新型溴代阻燃剂的水平及组成分布[J]. 环境科学, 2014, 35(4):1230-1237.
WU H, JIN J, WANG Y, et al. Comparative study of the level and distribution of polybrominated diphenyl ethers and new brominated flame retardants in the atmosphere of typical urban[J]. Environmental Science, 2014, 35(4):1230-1237 (in Chinese).
- [31] STRANDBERG B, DODDER N G, BASU I, et al. Concentrations and spatial variations of polybrominated diphenyl ethers and other organohalogen compounds in Great Lakes air[J]. Environmental Science & Technology, 2001, 35(6):1078-1083.
- [32] SU Y S, HUNG H, BRICE K A, et al. Air concentrations of polybrominated diphenyl ethers (PBDE) in 2002—2004 at a rural site in the Great Lakes[J]. Atmospheric Environment, 2009, 43(39):6230-6237.
- [33] GOUIN T, HAMER T, DALY G L. Variability of concentrations of polybrominated diphenyl ethers and polychlorinated biphenyls in air: Implications for monitoring, modeling and control[J]. Atmospheric Environment, 2005, 39(1):151-166.
- [34] HARRAD S, HUNTER S. Concentrations of polybrominated diphenyl ethers in air and soil on a rural-urban transect across a major UK conurbation[J]. Environmental Science & Technology, 2006, 40(15):4548-4553.
- [35] JULANDER A, WESTBERG H, ENGWALL M, et al. Distribution of brominated flame retardants in different dust fractions in air from an electronics recycling facility[J]. Science of the Total Environment, 2005, 350(1-3):151-160.
- [36] TIAN M, CHEN S J, WANG J, et al. Brominated flame retardants in the atmosphere of e-waste and rural sites in southern China: Seasonal variation, temperature dependence, and gas-particle partitioning[J]. Environmental Science & Technology, 2011, 45(20):8819-8825.
- [37] BUTT C M, DIAMOND M L, TRUONG J. Spatial distribution of polybrominated diphenyl ethers in southern Ontario as measured in indoor and outdoor window organic films[J]. Environmental Science & Technology, 2004, 38(3):724-731.
- [38] WILFORD B H, HARNER T, ZHU J P, et al. Passive sampling survey of polybrominated diphenyl ether flame retardant in indoor and outdoor air in Ottawa, Canada: implications for sources and exposure [J]. Environmental Science & Technology, 2004, 38(20):5312-5318.
- [39] CHEN L G, MAI B X, XU Z C, et al. In- and outdoor sources of polybrominated diphenyl ethers and their human inhalation exposure in Guangzhou, China[J]. Atmospheric Environment, 2008, 42(1):78-86.
- [40] TUE N M, TAKAHASHI S, SUZUKI G, et al. Contamination of indoor dust and air by polychlorinated biphenyls and brominated flame retardants and relevance of non-dietary exposure in Vietnamese informal e-waste recycling sites[J]. Environment International, 2013, 51(5):160-167.
- [41] LI Y, CHEN L, WEN Z H, et al. Characterizing distribution, sources, and potential health risk of polybrominated diphenyl ethers (PBDEs) in office environment[J]. Environmental Pollution, 2015, 198:25-31.
- [42] ZHUN Z, LIU L Y, MA W L, et al. Polybrominated diphenyl ethers (PBDEs) in the indoor dust in China: Levels, spatial distribution and human exposure[J]. Ecotoxicology and Environmental Safety, 2015, 111:1-8.
- [43] 韩文亮, 冯加良, 彭小霞, 等. 上海室内环境中多溴联苯醚的沉降通量与组成[J]. 环境科学, 2010, 31(3):579-585.
HAN W L, FENG J L, PENG X X. Indoor deposition flux and congener profiles of polybrominated diphenyl ethers (PBDEs) in Shanghai, China[J]. Environmental Science, 2010, 31(3):579-585 (in Chinese).
- [44] TOMY G, TITTELMIER U, BRAEKEVELT E, et al. The second international workshop on brominated flame retardants[C]. Stockholm: AB Firmatryck, 2001, 104-107.
- [45] WATANABE I, SAKAI S. Environmental release and behavior of brominated flame retardants[J]. Environment International, 2003, 29(6):665-682.
- [46] STREETS S S, HENDERSON S A, STONER A D, et al. Partitioning and bioaccumulation of PBDEs and PCBs in Lake Michigan[J]. Environmental Science & Technology, 2006, 40(23):7263-7269.
- [47] LAW K, HALLDORSON T, DANELL R, et al. Bioaccumulation and trophic transfer of some brominated flame retardants in a Lake Winnipeg (Canada) food web[J]. Environmental Toxicology and Chemistry, 2006, 25(8):2177-2186.
- [48] VECCHIATO M, ARGIRIADIS E, ZAMBON S, et al. Persistent organic pollutants (POPs) in Antarctica: Occurrence in continental and coastal surface snow[J]. Microchemical Journal, 2015, 119:75-82.
- [49] BOOIJ K, ZEGERS B N, BOON J P. Levels of some polybrominated diphenyl ether (PBDE) flame retardants along the Dutch coast as derived from their accumulation in SPMDs and blue mussels (*Mytilus edulis*) [J]. Chemosphere, 2002, 46(5):683-688.
- [50] LUO X J, YU M, MAI B X, et al. Distribution and partition of polybrominated diphenyl ethers (PBDEs) in water of the Zhujiang River Estuary[J]. Chinese Science Bulletin, 2008, 53(4):493-500.
- [51] BOGDAL C, SCHERINGER M, SCHMID P, et al. Levels, fluxes and time trends of persistent organic pollutants in Lake Thun, Switzerland: Combining trace analysis and multimedia modeling[J]. Science of the Total Environment, 2010, 408(17):3654-3663.
- [52] ORAM J J, MCKEE L J, WERME C E, et al. A mass budget of polybrominated diphenyl ethers in San Francisco Bay, CA [J]. Environment International, 2008, 34(8):1137-1147.
- [53] WANG J X, LIN Z K, LIN K F, et al. Polybrominated diphenyl ethers in water, sediment, soil, and biological samples from different industrial areas in Zhejiang, China[J]. Journal of Hazardous Materials, 2011, 197(6):211-219.
- [54] WU J P, LUO X J, ZHANG Y, et al. Bioaccumulation of polybrominated diphenyl ethers (PBDEs) and polychlorinated biphenyl (PCBs) in wild aquatic species from an electronic waste (e-waste) recycling site in South China [J]. Environment International, 2008, 34(8):

1109-1113.

- [55] GE J, LIU M X, YUN X Y, et al. Occurrence, distribution and seasonal variations of polychlorinated biphenyls and polybrominated diphenyl ethers in surface waters of the East Lake, China[J]. *Chemosphere*, 2014, 103(5):256-262.
- [56] SONG W, FORD J C, LI A, et al. Polybrominated diphenyl ethers in the sediments of the Great Lakes. 1. Lake Superior [J]. *Environmental Science & Technology*, 2004, 38(12):3286-3293.
- [57] SONG W, LI A, FORD J C, et al. Polybrominated diphenyl ethers in the sediments of the Great Lakes. 2. Lakes Michigan and Huron[J]. *Environmental Science & Technology*, 2005, 39(10):3474-3479.
- [58] SONG W, FORD J C, LI A, et al. Polybrominated diphenyl ethers in the sediments of the Great Lakes. 3. Lake Ontario and Erie[J]. *Environmental Science & Technology*, 2005, 39(15):5600-5605.
- [59] ZHU H W, WANG Y, WANG X W, et al. Distribution and accumulation of polybrominated diphenyl ethers (PBDEs) in Hong Kong mangrove sediments[J]. *Science of the Total Environment*, 2014, 468-469:130-139.
- [60] MOON H B, KANNAN K, CHOI M, et al. Polybrominated diphenyl ethers (PBDEs) in marine sediments from industrialized bays of Korea [J]. *Marine Pollution Bulletin*, 2007, 54(9):1402-1412.
- [61] MOON H B, KANNAN K, LEE S J, et al. Polybrominated diphenyl ethers (PBDEs) in sediment and bivalves from Korean coastal waters [J]. *Chemosphere*, 2007, 66(2):243-251.
- [62] BRANDSMA S H, LEONARDS P E, LESLIE H A, et al. Tracing organophosphorus and brominated flame retardants and plasticizers in an estuarine food web[J]. *Science of the Total Environment*, 2015, 505(505C):22-31.
- [63] MAI B X, CHEN S J, LUO X J, et al. Distribution of polybrominated diphenyl ethers in sediments of the Pearl River Delta and adjacent South China Sea[J]. *Environmental Science & Technology*, 2005, 39(10):3521-3527.
- [64] CHEN S J, GAO X J, MAI B X, et al. Polybrominated diphenyl ethers in surface sediments of the Yangtze River Delta: Levels, distribution and potential hydrodynamic influence[J]. *Environmental Pollution*, 2006, 144(3):951-957.
- [65] 吕建霞,王亚韡,张庆华,等.天津大沽排污河口沉积物多溴联苯醚、有机氯农药和重金属的污染趋势[J].*科学通报*,2007,52(3):277-282.
- LÜ J X, WANG Y W, ZHANG Q H. Contamination trends of polybrominated diphenyl ethers, organochlorine pesticides and heavy metals in sediments from Dagu Drainage River estuary, Tianjin[J]. *Chinese Science Bulletin*, 2007, 52(3):277-282 (in Chinese).
- [66] JIANG J J, LEE C L, FANG M D, et al. Polybrominated diphenyl ethers and polychlorinated biphenyls in sediments of southwest Taiwan: Regional characteristics and potential sources[J]. *Marine Pollution Bulletin*, 2011, 62(4):815-823.
- [67] WANG T, LI S S, ZHANG C, et al. Quantities, sources and adsorption of polybrominated diphenyl ethers in components of surficial sediments collected in Songhua River (Jilin City), China[J]. *Chemosphere*, 2015, 119(119C):1208-1216.
- [68] LIU L L, LI H, WANG Z P, et al. Insights into spatially and temporally co-occurring polybrominated diphenyl ethers in sediments of the East China Sea[J]. *Chemosphere*, 2015, 123:55-63.
- [69] MARVIN C, WILLIAMS D, KUNTZ K, et al. Temporal trends in polychlorinated dibenzo-*p*-dioxins and dibenzofurans, dioxin-like PCBs, and polybrominated diphenyl ethers in Niagara river suspended sediments[J]. *Chemosphere*, 2007, 67(9):1808-1815.
- [70] NORTH K D. Tracking polybrominated diphenyl ether releases in a wastewater treatment plant effluent, Palo Alto, California [J]. *Environmental Science & Technology*, 2004, 38(17):4484-4488.
- [71] KNOTH W, MANN W, MEYER R, et al. Polybrominated diphenyl ether in sewage sludge in Germany[J]. *Chemosphere*, 2007, 67(9):1831-1837.
- [72] WANG Y W, ZHANG Q H, LV J X, et al. Polybrominated diphenyl ethers and organochlorine pesticides in sewage sludge of wastewater treatment plants in China[J]. *Chemosphere*, 2007, 68(9):1683-1691.
- [73] ELJARRAT E, MARSH G, LABANDEIRA A, et al. Effect of sewage sludges contaminated with polybrominated diphenyl ethers on agricultural soils[J]. *Chemosphere*, 2008, 71(6):1079-1086.
- [74] GEVAO B, MUZAINI S, HELALEH M, et al. Occurrence and concentrations of polybrominated diphenyl ethers in sewage sludge from three wastewater treatment plants in Kuwait[J]. *Chemosphere*, 2008, 71(2):242-247.
- [75] ANDRADE N A, MCCONNELL L L, TORRENTS A, et al. Persistence of polybrominated diphenyl ethers in agricultural soils after biosolids applications[J]. *Journal of Agricultural & Food Chemistry*, 2010, 58(5):3077-3084.
- [76] PENG X Z, TANG C M, YU Y Y, et al. Concentrations, transport, fate, and releases of polybrominated diphenyl ethers in sewage treatment plants in the Pearl River Delta, South China[J]. *Environment International*, 2009, 35(2):303-309.
- [77] STIBOROVA H, VRKOSLAVOVA J, LOVECKA P, et al. Aerobic biodegradation of selected polybrominated diphenyl ethers (PBDEs) in wastewater sewage sludge[J]. *Chemosphere*, 2015, 118:315-321.
- [78] HASSANIN A, BREIVIK K, MEIJER S N, et al. PBDEs in European background soils: levels and factors controlling their distribution [J]. *Environmental Science & Technology*, 2004, 38(3):738-745.
- [79] LEUNG A O, LUKSEMBURG W J, WONG A S, et al. Spatial distribution of polybrominated diphenyl ethers and polychlorinated dibenzo-*p*-dioxins and dibenzofurans in soil and combusted residue at Guiyu, an electronic waste recycling site in Southeast China [J]. *Environmental Science & Technology*, 2007, 41(8):2730-2737.

- [80] WANG P, ZHANG Q H, WANG Y W, et al. Altitude dependence of polychlorinated biphenyls (PCBs) and polybrominated diphenyl ethers (PBDEs) in surface soil from Tibetan Plateau, China[J]. *Chemosphere*, 2009, 76(11):1498-1504.
- [81] MA J, ADDINK R, YUN S, et al. Polybrominated dibenzo-*p*-dioxins/dibenzofurans and polybrominated diphenyl ethers in soil, vegetation, workshop-floor dust, and electronic shredder residue from an electronic waste recycling facility and in soils from a chemical industrial complex in Eastern China[J]. *Environmental Science & Technology*, 2009, 43(19):7350-7356.
- [82] WANG Y, LUO C L, LI J, et al. Characterization of PBDEs in soils and vegetations near an e-waste recycling site in South China[J]. *Environmental Pollution*, 2011, 159(10):2443-2448.
- [83] ZHENG X Y, LIU X D, JIANG G B, et al. Distribution of PCBs and PBDEs in soils along the altitudinal gradients of Balang Mountain, the east edge of the Tibetan Plateau[J]. *Environmental Pollution*, 2012, 161(1):101-106.
- [84] YUAN G L, HAN P, XIE W, et al. Altitudinal distribution of polybrominated diphenyl ethers (PBDEs) in the soil along Central Tibetan Plateau, China[J]. *Science of the Total Environment*, 2012, 433(7):44-49.
- [85] SUN Y, YUAN G L, LI J, et al. Polybrominated diphenyl ethers (PBDEs) in surface soils near the Changwengluozha Glacier of Central Tibetan Plateau, China[J]. *Science of the Total Environment*, 2015, 511:399-406.
- [86] CHENG Z N, WANG Y, WANG S R, et al. The influence of land use on the concentration and vertical distribution of PBDEs in soils of an e-waste recycling region of South China[J]. *Environmental Pollution*, 2014, 191:126-131.
- [87] NIE Z Q, TIAN S L, TIAN Y J, et al. The distribution and biomagnification of higher brominated BDEs in terrestrial organisms affected by a typical e-waste burning site in South China[J]. *Chemosphere*, 2015, 118C(1):301-308.
- [88] ZHENG X B, LUO X J, ZHENG J, et al. Contaminant sources, gastrointestinal absorption, and tissue distribution of organohalogenated pollutants in chicken from an e-waste site[J]. *Science of the Total Environment*, 2015, 505(505C):1003-1010.
- [89] 高永飞,王璞,陈卫海,等. 喜马拉雅山区莱籽草和棘豆样品中 PCBs, PBDEs 和 PCDD/Fs 的分析[J]. *环境化学*, 2010, 31(1):26-30.
- GAO Y F, WANG P, CHEN W H, et al. PCBs, PBDEs and PCDD/Fs in the grass samples from the Himalayas[J]. *Environmental Chemistry*, 2010, 31(1):26-30 (in Chinese).
- [90] MAHMOOD A, MALIK R N, SYED J H, et al. Dietary exposure and screening-level risk assessment of polybrominated diphenyl ethers (PBDEs) and dechloran plus (DP) in wheat, rice, soil, and air along two tributaries of the River Chenab, Pakistan[J]. *Chemosphere*, 2015, 118(1):57-64.
- [91] JIN J, LIU W Z, WANG Y, et al. Levels and distribution of polybrominated diphenyl ethers in plant, shellfish and sediment samples from Laizhou Bay in China[J]. *Chemosphere*, 2008, 71(6):1043-1050.
- [92] SALAMOVA A, HITES R A. Brominated and chlorinated flame retardants in tree bark from around the globe[J]. *Environmental Science & Technology*, 2013, 47(1):349-354.
- [93] KELLY B C, IKONOMOU M G, BLAIR J D, et al. Bioaccumulation behaviour of polybrominated diphenyl ethers (PBDEs) in a Canadian Arctic marine food web[J]. *Science of The Total Environment*, 2008, 401(1-3):60-72.
- [94] 史双昕,曾良子,周丽,等. 江苏经济高速发展城市香樟树皮中的多溴联苯醚[J]. *环境科学*, 2011, 32(9):2654-2660.
- SHI S X, ZENG L Z, ZHOU L, et al. Polybrominated diphenyl ethers in camphor bark from speedy developing urban in Jiangsu Province [J]. *Environmental Science*, 2011, 32(9):2654-2660 (in Chinese).
- [95] ZHAO Y L, YANG L M, WANG Q Q. Modeling persistent organic pollutant (POP) partitioning between tree bark and air and its application to spatial monitoring of atmospheric POPs in mainland China[J]. *Environmental Science & Technology*, 2008, 42(16):6046-6051.
- [96] YUAN H D, JIN J, BAI Y, et al. Concentrations and distributions of polybrominated diphenyl ethers and novel brominated flame retardants in tree bark and human hair from Yunnan Province, China[J]. *Chemosphere*, 2016, 154:319-325.
- [97] ZHU L, HITES R A. Brominated flame retardants in tree bark from North America[J]. *Environmental Science & Technology*, 2006, 40(12):3711-3716
- [98] WEN S, YANG F, LI J G, et al. Polychlorinated dibenzo-*p*-dioxin and dibenzofurans (PCDD/Fs), polybrominated diphenyl ethers (PBDEs), and polychlorinated biphenyls (PCBs) monitored by tree bark in an E-waste recycling area[J]. *Chemosphere*, 2009, 74(7):981-987.
- [99] HU J C, JIN J, WANG Y, et al. Levels of polybrominated diphenyl ethers and hexabromocyclododecane in the atmosphere and tree bark from Beijing, China[J]. *Chemosphere*, 2011, 84(3):355-360.
- [100] WANG S, ZHANG S Z, HUANG H L, et al. Characterization of polybrominated diphenyl ethers (PBDEs) and hydroxylated and methoxylated PBDEs in soils and plants from an e-waste area, China[J]. *Environmental Pollution*, 2014, 184:405-413.
- [101] DOMINGO J L, MARTI-CID R, CASTELL V, et al. Human exposure to PBDEs through the diet in Catalonia, Spain: Temporal trend. A review of recent literature on dietary PBDE intake[J]. *Toxicology*, 2008, 248(1):25-32.
- [102] ZHU C F, LI Y M, WANG P, et al. Polychlorinated biphenyls (PCBs) and polybrominated biphenyl ethers (PBDEs) in environmental samples from Ny-Ålesund and London Island, Svalbard, the Arctic[J]. *Chemosphere*, 2015, 126(1):40-46.
- [103] ST-AMAND A D, MAYER P M, BLAIS J M. Seasonal trends in vegetation and atmospheric concentrations of PAHs and PBDEs near a

- sanitary landfill[J]. *Atmospheric Environment*, 2008, 42(13):2948-2958.
- [104] 刘娟,罗孝俊,陈社军,等. 电子垃圾回收地散养鸡(鸡)中多溴联苯醚(PBDEs)的浓度与分布模式[J]. *现代农业科技*, 2009, (24):293-297.
- LIU J, LUO J X, CHEN S J, et al. Concentration and distribution pattern of polybrominated diphenyl ethers (PBDEs) in scattered Chicken from electronic waster recycling site[J]. *Modern Agricultural Science and Technology*, 2009, (24):293-297(in Chinese).
- [105] 胡国成,许振成,戴家银,等. 有机氯农药和多溴联苯醚在白洋淀鸭子组织中分布特征研究[J]. *环境科学*, 2010, 31(12):3081-3087.
- HU G C, XU Z C, DAI J Y, et al. Distribution characteristic of organochlorine pesticides and polybrominated diphenyl ethers in tissues of ducks from Baiyangdian Lake, North China[J]. *Environmental Science*, 2010, 31(12):3081-3087(in Chinese).
- [106] QIN X F, QIN Z F, LI Y, et al. Polybrominated diphenyl ethers in chicken tissues and eggs from an electronic waste recycling area in southeast China[J]. *Journal of Environmental Sciences*, 2011, 23(1):133-138.
- [107] TANG W B, HUANG K, ZHAO J H, et al. Polybrominated diphenyl ethers in resident Eurasian Tree Sparrow from Shanghai: Geographical distribution and implication for potential sources[J]. *Chemosphere*, 2015, 126:25-31.
- [108] HUANG K, LIN K F, GUO J, et al. Polybrominated diphenyl ethers in birds from Chongming Island, Yangtze estuary, China: Insight into migratory behavior[J]. *Chemosphere*, 2013, 91(10):1416-1425.
- [109] DASO A P, OKONKWO J O, JANSEN R, et al. Polybrominated diphenyl ethers (PBDEs) in eggshells of the Southern Ground-Hornbill (*Bucorvus leadbeateri*) and Wattled Crane (*Bugeranus carunculatus*) in South Africa[J]. *Chemosphere*, 2015, 118(1):284-292.
- [110] HU G C, LUO X J, DAI J Y, et al. Brominated flame retardants, polychlorinated biphenyls, and organochlorine pesticides in captive giant panda (*Ailuropoda melanoleuca*) and red panda (*Ailurus fulgens*) from China[J]. *Environmental Science & Technology*, 2008, 42(13):4704-4709.
- [111] KANNAN K, YUN S H, EVANS T J. Chlorinated, brominated, and perfluorinated contaminants in livers of polar bears from Alaska[J]. *Environmental Science & Technology*, 2005, 39(23):9057-9063.
- [112] YU L H, LUO X J, WU J P, et al. Biomagnification of higher brominated PBDE congeners in an urban terrestrial food web in North China based on field observation of prey deliveries[J]. *Environmental Science & Technology*, 2011, 45(12):5125-5131.
- [113] YANG R Q, JING C Y, ZHANG Q H, et al. Polybrominated diphenyl ethers (PBDEs) and mercury in fish from lakes of the Tibetan Plateau[J]. *Chemosphere*, 2011, 83(6):862-867.
- [114] MIÈGE C, PERETTI A, LABADIE P, et al. Occurrence of priority and emerging organic compounds in fishes from the Rhone River (France)[J]. *Analytical and Bioanalytical Chemistry*, 2012, 404(9):2721-2735.
- [115] JÜRGENS M D, JOHNSON A C, JONES K C, et al. The presence of EU priority substances mercury, hexachlorobenzene, hexachlorobutadiene and PBDEs in wild fish from four English rivers[J]. *Science of the Total Environment*, 2013, 461-462(7):441-452.
- [116] SUN R X, LUO X J, TANG B, et al. Persistent halogenated compounds in fish from rivers in the Pearl River Delta, South China: Geographical pattern and implications for anthropogenic effects on the environment[J]. *Environmental Research*, 2016, 146(6):371-378.
- [117] 郭英,唐洪磊,孟祥周,等. 多溴联苯醚在桂花鱼体内的分布[J]. *环境科学*, 2007, 28(12):2806-2810.
- GUO Y, TANG H L, MENG X Z, et al. Tissue distribution of PBDEs in Mandarin fish[J]. *Environmental Science*, 2007, 28(12):2806-2810(in Chinese).
- [118] 向彩红,罗孝俊,余梅,等. 珠江河口水生生物中多溴联苯醚的分布[J]. *环境科学*, 2006, 27(9):1732-1737.
- XIANG C H, LUO X J, YU M, et al. Distribution of polybrominated diphenyl ethers in aquatic species from the Pearl River estuary[J]. *Environmental Science*, 2006, 27(9):1732-1737(in Chinese).
- [119] 王俊霞,王春艳,刘莉莉,等. 多溴联苯醚在市场鲫鱼体内分布和食鱼暴露量[J]. *环境科学*, 2014, 35(8):3175-3182.
- WANG J X, WANG C Y, LIU L L, et al. Distribution of polybrominated diphenyl ethers in wild crucian carp and exposure estimation of dietary intake[J]. *Environmental Science*, 2014, 35(8):3175-3182(in Chinese).
- [120] 田胜艳,梁贤伟,祝凌燕. 多溴联苯醚的生物富集与生物有效性//第六届全国环境化学大会暨环境科学仪器与分析仪器展览会摘要集[C]. 2011年.
- TIAN S Y, LIANG X W, ZHU L Y. Bioaccumulation and bioavailability of polybrominated diphenyl ethers//The abstract book of the 6th national conference on environmental chemistry and the exhibition of environmental science instruments and analytical instruments[C]. 2011(in Chinese).
- [121] SHE J W, PETREAS M, WINKLER J, et al. PBDEs in the San Francisco Bay area: Measurements in harbor seal blubber and human breast adipose tissue[J]. *Chemosphere*, 2002, 46(5):697-707.
- [122] RAMU K, KAJIWARA N, LAM P K S, et al. Temporal variation and biomagnification of organohalogen compounds in finless porpoises (*Neophocaena phocaenoides*) from the South China Sea[J]. *Environmental Pollution*, 2006, 144:516-523.
- [123] ROTANDER A, BAVEL B V, POLDER A, et al. Polybrominated diphenyl ethers (PBDEs) in marine mammals from Arctic and North Atlantic regions, 1986-2009[J]. *Environment International*, 2012, 40(2):102-109.
- [124] ROSS P S, NOËL M, LAMBOUM D, et al. Declining concentrations of persistent PCBs, PBDEs, PCDEs, and PCNs in harbor seals

- (*Phoca vitulina*) from the Salish Sea[J]. *Progress in Oceanography*, 2013, 115(4):160-170.
- [125] HOUDE M, WANG X, FERGUSON S H, et al. Spatial and temporal trends of alternative flame retardants and polybrominated diphenyl ethers in ringed seals (*Phoca hispida*) across the Canadian Arctic[J]. *Environmental Pollution*, 2017, 223:266-276.
- [126] SHI Z X, JIAO Y, HU Y, et al. Levels of tetrabromobisphenol A, hexabromocyclododecanes and polybrominated diphenyl ethers in human milk from the general population in Beijing, China[J]. *Science of the Total Environment*, 2013, 452-453(452-453C):10-18.
- [127] PARK J S, SHE J W, HOLDEN A, et al. Postnatal exposures to polybrominated diphenyl ethers (PBDEs) and polychlorinated biphenyls (PCBs) via breast milk in California: Does BDE-209 transfer to breast milk? [J]. *Environmental Science & Technology*, 2011, 45(10):4579-4585.
- [128] ZHU L Y, MA B L, LI J G, et al. Distribution of polybrominated diphenyl ethers in breast milk from North China: Implication of exposure pathways[J]. *Chemosphere*, 2009, 74(11):1429-1434.
- [129] CHOVANCOVÁ J, ČONKA K, KOČAN A, et al. PCDD, PCDF, PCB and PBDE concentrations in breast milk of mothers residing in selected areas of Slovakia[J]. *Chemosphere*, 2011, 83(10):1383-1390.
- [130] TOMS L M, HARDEN F A, SYMONS R K, et al. Polybrominated diphenyl ethers (PBDEs) in human milk from Australia[J]. *Chemosphere*, 2007, 68(5):797-803.
- [131] FUJII Y, ITO Y, HARADA K H, et al. Regional variation and possible sources of brominated contaminants in breast milk from Japan[J]. *Environmental Pollution*, 2012, 162:269-274.
- [132] MA S T, YU Z Q, ZHANG X L, et al. Levels and congener profiles of polybrominated diphenyl ethers (PBDEs) in breast milk from Shanghai: Implication for exposure route of higher brominated BDEs[J]. *Environment International*, 2012, 42(1):72-77.
- [133] DIMITRIADOU L, MALARVANNAN G, COVACI A, et al. Levels and profiles of brominated and chlorinated contaminants in human breast milk from Thessaloniki, Greece[J]. *Science of the Total Environment*, 2016, 539(1):350-358.
- [134] LEE S, KIM S, KIM E, et al. Polybrominated diphenyl ethers (PBDEs) in breast milk of Korea in 2011: Current contamination, time course variation, influencing factors and health risks[J]. *Environmental Research*, 2013, 126(4):76-83.
- [135] GÓMARA B, HERRERO L, PACEPAVICIUS G, et al. Occurrence of co-planar polybrominated/chlorinated biphenyls (PXBs), polybrominated diphenyl ethers (PBDEs) and polychlorinated biphenyls (PCBs) in breast milk of women from Spain[J]. *Chemosphere*, 2011, 83:799-805.
- [136] MEIRONYTĖ D, NORÉN K, BERGMAN Å. Analysis of polybrominated diphenyl ethers in Swedish human milk. A time-related trend study, 1972-1997[J]. *Journal of Toxicology and Environmental Health*, 1999, 58(6):329-341.
- [137] FROMME H, ALBRECHT M, APPEL M, et al. PCBs, PCDD/Fs, and PBDEs in blood samples of a rural population in South Germany [J]. *International Journal of Hygiene and Environmental Health*, 2015, 218(1):41-46.
- [138] KIM J, SON M H, SHIN E S et al. Occurrence of dechlorane compounds and polybrominated diphenylethers (PBDEs) in the Korean general population[J]. *Environmental Pollution*, 2016, 212:330-336.
- [139] UEMURA H, ARISAWA K, HIYOSHI M, et al. Congener-specific body burden levels and possible determinants of polybrominated diphenyl ethers in the general Japanese population[J]. *Chemosphere*, 2010, 79(7):706-712.
- [140] SHEN H T, DING G Q, HAN G G, et al. Distribution of PCDD/Fs, PCBs, PBDEs and organochlorine residues in children's blood from Zhejiang, China[J]. *Chemosphere*, 2010, 80(2):170-175.
- [141] ZHU L Y, MA B L, HITES R A. Brominated flame retardants in serum from the general population in northern China[J]. *Environmental Science & Technology*, 2009, 43(18):6963-6968.
- [142] 屈伟月,王德超,盛国英,等. 婴儿脐带血和母亲血中多溴联苯醚的研究[J]. *中国环境科学*, 2007, 27(2):269-272.
- QU W Y, WANG D C, SHENG G Y, et al. Preliminary studies on polybrominated diphenyl ethers in fetal and maternal blood[J]. *China Environmental Science*, 2007, 27(2):269-272(in Chinese).
- [143] STASINSKA A, HEYWORTH J, REID A, et al. Polybrominated diphenyl ether (PBDE) concentrations in plasma of pregnant women from Western Australia[J]. *Science of the Total Environment*, 2014, 493(5):554-561.
- [144] MA W L, YUN S H, BELL E M, et al. Temporal trends of polybrominated diphenyl ethers (PBDEs) in the blood of newborns from New York state during 1997 through 2011: Analysis of dried blood spots from the newborn screening program[J]. *Environmental Science & Technology*, 2013, 47(14):8015-8021.
- [145] KRÓL S, NAMIEŚNIK J, ZABIEGALA B. Occurrence and levels of polybrominated diphenyl ethers (PBDEs) in house dust and hair samples from Northern Poland; an assessment of human exposure[J]. *Chemosphere*, 2014, 110:91-96.
- [146] ZHENG J, LUO X J, YUAN J G, et al. Levels and sources of brominated flame retardants in human hair from urban, e-waste, and rural areas in South China[J]. *Environmental Pollution*, 2011, 159(12):3706-3713.
- [147] LIU P Y, ZHAO Y X, ZHU Y Y, et al. Determination of polybrominated diphenyl ethers in human semen[J]. *Environment International*, 2012, 42(1):132-137.
- [148] LEONETTI C, BUTT C M, HOFFMAN K, et al. Concentrations of polybrominated diphenyl ethers (PBDEs) and 2,4,6-tribromophenol in human placental tissues[J]. *Environment International*, 2016, 88:23-29.
- [149] 张茨,吴江平,罗孝俊,等. 多溴联苯醚在典型电子垃圾污染区域水生食物链上的生物富集特征[J]. *生态毒理学报*, 2009, 4(3):

338-344.

ZHANG Y, WU J P, LUO X J, et al. Bioaccumulation of polybrominated diphenyl ethers in a contaminated freshwater food web from an E-waste recycling site, South China[J]. *Asian Journal of Ecotoxicology*, 2009, 4(3):338-344(in Chinese).

- [150] DI PAOLO C, GANDHI N, BHAUSAR S P, et al. Black carbon inclusive multichemical modeling of PBDE and PCB biomagnification and transformation in estuarine food webs[J]. *Environmental Science & Technology*, 2010, 44(19):7548-7554.
- [151] KOBAYASHI J, IMUTA Y, KOMORITA T, et al. Trophic magnification of polychlorinated biphenyls and polybrominated diphenyl ethers in an estuarine food web of the Ariake Sea, Japan[J]. *Chemosphere*, 2015, 118:201-206.
- [152] MA X D, ZHANG H J, YAO Z W, et al. Bioaccumulation and trophic transfer of polybrominated diphenyl ethers(PBDEs) in a marine food web from Liaodong Bay, North China[J]. *Marine Pollution Bulletin*, 2013, 74(1):110-115.
- [153] SHAW S D, BERGER M L, BRENNER D, et al. Bioaccumulation of polybrominated diphenyl ethers and hexabromocyclododecane in the northwest Atlantic marine food web[J]. *Science of the Total Environment*, 2009, 407(10):3323-3329.
- [154] MUIR D C, BACKUS S, DEROCHE A E, et al. Brominated flame retardants in polar bears (*Ursus maritimus*) from Alaska, the Canadian Arctic, East Greenland, and Svalbard[J]. *Environmental Science & Technology*, 2006, 40(2):449-455.
- [155] VOORSPOELS S, COVACI A, JASPERS V L, et al. Biomagnification of PBDEs in three small terrestrial food chains[J]. *Environmental Science & Technology*, 2007, 41(2):411-416.
- [156] ZHENG B H, ZHAO X R, NI X J, et al. Bioaccumulation characteristics of polybrominated diphenyl ethers in the marine food web of Bohai Bay[J]. *Chemosphere*, 2016, 150:424-430.
- [157] SHAO M H, TAO P, WANG M, et al. Trophic magnification of polybrominated diphenyl ethers in the marine food web from coastal area of Bohai Bay, North China[J]. *Environmental Pollution*, 2016, 213:379-385.
- [158] 吴江平,张炎,罗孝俊,等. 多溴联苯醚的生物富集效应研究进展[J]. *生态毒理学报*, 2009, 4(2):153-163.
WU J P, ZHANG Y, LUO J X, et al. A review on the bioaccumulation behavior of polybrominated diphenyl ethers[J]. *Asian Journal of Ecotoxicology*, 2009, 4(2):153-163(in Chinese).
- [159] BRAGIGAND V, AMIARD-TRIQUET C, PARLIER E, et al. Influence of biological and ecological factors on the bioaccumulation of polybrominated diphenyl ethers in aquatic food webs from French estuaries[J]. *Science of the Total Environment*, 2006, 368(2-3):615-626.